



International Journal of Advanced Academic Studies

E-ISSN: 2706-8927

P-ISSN: 2706-8919

www.allstudyjournal.com

IJAAS 2023; 5(11): 21-24

Received: 21-09-2023

Accepted: 28-10-2023

Himanshi

Researcher, Department of
Food Technology, Desh
Bhagat University Punjab,
India

Dr. Ritu Aggarwal

Major Advisor, Department of
Food Technology, Desh
Bhagat University Punjab,
India

Deepak

Assistant Professor and
Advisor, Department of
Agriculture and Technology,
Maharana Pratap University,
Udaipur, Rajasthan, India

Corresponding Author:

Himanshi

Researcher, Department of
Food Technology, Desh
Bhagat University Punjab,
India

Cookies enriched with protein manufactured with (Soya Flour) and Fibre (Banana Peel Powder)

Himanshi, Dr. Ritu Aggarwal and Deepak

DOI: <https://doi.org/10.33545/27068919.2023.v5.i11a.1076>

Abstract

This study aimed to evaluate the effect of incorporation of banana peel flour (BPF) on the nutritional and antioxidant properties of cookies prepared with wheat flour. Cookies were prepared by substituting 7.5, 10, 12.5, and 15% of superfine wheat flour with BPF and compared to control samples (with no BPF). The results showed significant differences in the moisture, protein, crude fat, and ash contents of 3.9-7.5, 1.75-8.75, 12.3-25.9, and 1.3-2.00 g/100 g, respectively. The highest moisture and ash contents were recorded in cookies with 15% BPF, while the highest protein and crude fat contents were recorded in control cookies. Physical characteristics such as the weight, diameter, thickness, and spread ratio were found to be 8.87-12.49 g, 4.067-4.167 cm, 0.4-1.13 cm, and 3.683-10.587, respectively. The hardness of cookies was increased significantly with the increasing proportion of BPF in cookies at a range of 15.84-25.55 N. However, the hardness of the cookies decreased with storage. Color range values, L^* , a^* , and b^* , of the cookies were found to be 38.85-52.81, 6.74-8.12, and 16.09-21.52, respectively, with the higher L^* , a^* , and b^* values for control cookies. However, these values changed during storage. Moreover, antioxidant properties significantly increased in cookies with the substitution of BPF, where the total phenolic content, DPPH, and lipid peroxidation were found to be in the range of 0.282-0.921 mg GAE/g, 47.761-70.298, and 36.74-71.84%, respectively, and some values decreased during storage. The results showed that cookies enriched with BPF were obtained with improved antioxidant properties and total phenolic content without altering their physical and nutritional properties and also fathomed the favorable acceptance.

Keywords: Banana peel flour, cookies and storage

Introductions

The bakery industry in India plays an important role within the industrial map of this country. Bakery products are things of mass consumption because of its low pricing and palatability. Due to moving eating habit of consumer growth of bakery industry has been significantly noticed. This industry, typically, consist of cookies, biscuits, breads cakes etc. Among the bakery products cookies and biscuits constitute to the most popular group.

Cookie is defined as a small thin crisp cake made from unleavened dough. Cookie are an important baked product in human diet and are usually eaten with tea and are also used as weaning food for infants. Cookie are ideal for their nutritive value, palatability, compactness and convenience. Having low moisture content than cakes and bread, Cookies are generally safer from microbiological spoilage and have long shelf-life, Akubor.

Cookies can be classified as ready to eat and convenient foods. Traditionally, the process of cookies making are fairly simple with basic ingredients consist of flour, eggs and sugar. Generally, cookies are recognized as flat, hard and crunchy food. Normally, cookies are classified according to their method of preparation such as drop, molded, presses, refrigerated, bar or rolled. Apart from that, the dominant ingredients that been used in the formulation also commonly being used to classify the cookies, for example, nut cookies, fruit cookies and chocolate cookies, Norhidayah 2014^[4].

Cookie is the product consumed from infant to old and now a day's because of health awareness they demanded fortified and enriched products with high nutritional value. Development of fortified cookie or other composite flour bakery products is the latest trend in the bakery industry. Most of bakery products are used as a source of incorporation of different nutritionally rich ingredient for their diversification. With increasing public health awareness worldwide, demand for functional food with multiple health benefits has also increased.

The use of medicinal food from folk medicine to prevent diseases such as diabetes, obesity, and cardiovascular problems is now gaining momentum among the public. Consumers are demanding foods that shows two main properties: the first-one deals with traditional nutritional aspects of the food, whereas, second feature, additional benefits are expected from its regular ingestion. These kinds of food products are often called functional foods. Varunkumar Here we are making efforts to raise the nutritive value of Cookies by replacing Refined Wheat flour (RWF) with Soya flour and Banana peel powder (BPP).

Main Ingredients

Soya bean

Soybean (*Glycine max* L.) is a species of legume native to East Asia, widely grown for its edible bean which has several uses.

The soybean is a valuable legume because it does provide all of the essential amino acids for humans; however it is relatively low in the sulfur containing amino acids, cysteine and methionine. It is one of the few legumes that can be consumed as a complete protein.

The soybean is comprised of approximately 37-42% protein.

Banana peel

The banana peel is a household and industrial food waste discarded in large quantities in nature. It represents about 35% of the total fresh mass of ripe fruit and there is not further involved in remarkable industrial applications. Bananas are one of the fruits most produced and consumed worldwide and the potential use of the peel would be of great relevance. Some researchers have revealed that the banana peel has compounds and nutrients important for food and for food industry. The banana peel is rich in dietary fiber, protein, essential amino acids, polyunsaturated fatty acids and potassium. It also contains antioxidant compounds including polyphenols, catecholamines and carotenoids, Rebello 2014, Spiller Mohammad A. ^[11] suggested that soy protein can be used as a source of high-quality protein to help satisfy the higher need for protein during muscle-building by providing the necessary essential amino acids for Wachirasir ^[17] shows that important constituent present in banana peels it also suggests. The main purpose of using banana peel is its fiber content. It contents around 50.25±0.2% of Total dietary fibers, Dietary fiber has shown beneficial effects in the prevention of several disease, such as cardiovascular diseases, diverticulosis, constipation, colon cancer and diabetes. The fruit fiber has a Better quality than other fiber sources due to its high total and soluble fiber content, water and oil holding capacities, fermentation ability, as well as a lower phytic acid and caloric value content.

Materials and Methods

This chapter includes the materials required and method adopted in the study. Present study on Development of cookies enriched with protein and fiber was carried out at Laboratory, University Department of Food Technology, Maharana Pratap University of Agricultural and technology, Udaipur. The banana peel was collected from the local manufacturer of banana chips. The major ingredients for the preparation of products were RWF, Soya flour, Powdered Sugar, Butter, Milk Powder and other chemicals were used from the laboratory store and Meera chemicals, Udaipur.

Raw materials

1. Refined Wheat flour

2. Soya flour
3. Banana Peel Powder
4. Butter
5. Milk Powder
6. Sugar
7. Egg

Chemicals

1. Baking Powder
2. Sodium bicarbonate
3. Essence

Proximate analysis

Moisture Content: The moisture content was determined as per standard methods (AOAC, 2000 ^[2]) and results were expressed in terms of percentage. The moisture content of as it is practiced in lab. Calibrating the weight of the Petri dish, than weigh of the sample 5gm was taken. The sample along with the petri dish was placed in the oven for 3 hours and maintained at the temperature at 105 °C, by repeating the process of drying, cooling and weighed at 30 min interval until it become to a constant weight. Then it was transferred to desiccators and weight of each sample as soon as the dishes are cool. Weight losses were calculated in each samples and get the average moisture content of sample. The percent moisture content was calculated as:

$$\text{Moisture (\%)} = \frac{\text{Weight loss}}{\text{Weight of the sample}} * 100$$

$$\text{Total solid \%} = 100 - \text{moisture.}$$

Ash content: The ash content was determined with the help of muffle furnace as per standard methods (AOAC, 2000 ^[2]) and results were expressed in terms of percentage. The ash content is a measure of the total amount of minerals present within a food, whereas the mineral content is a measure of the amount of specific inorganic components present within a food, such as Ca, Na, K and CL. Water and other volatile materials are vaporized and organic substances are burned in the presence of the oxygen in air to CO₂, H₂O and N₂. Ash contents of fresh foods rarely exceed 5%, although some processed foods can have ash contents as high as 12%. Some necessary apparatus include:

- Crucible (or similar porcelain or metal dishes)
- Muffled furnace
- Hot plate

To perform Ash content analysis product sample is powdered and weight is measured up to 110gm, after weight of the crucible is measured. Required amount of sample is introduced in crucible and the crucible is place in Muffle furnace at 550-600 °C for 4 to 5 hrs. The food sample is weighed before and after ashing to determine the concentration of ash present. The ash content can be expressed on either a *dry* or *wet* basis:

$$\% \text{ ash (Dry basis)} = \frac{\text{Weight after ashing} \div \text{Weight before ashing}}{\text{Weight before ashing}} * 100$$

Protein content: The protein content was determined with the help of Kjeldahl apparatus method as per the standard method (AACC, 2006 ^[1]) and results were expressed in terms of percentage. 0.5 gm. of ground sample by digesting with concentrated sulfuric acid at 100 °C. Then it was distilled with 40 per cent NaOH liberated ammonia was

trapped in 4% boric acid, using mix indicator (Methyl Red: bromocresol green 1:5). Then titrate it with 0.1NHCL, the percent of nitrogen was estimated and protein percentage was calculated by multiplying percent nitrogen with factor 6.25 Calculation:

$$\% \text{ nitrogen} = \frac{(S-B) \text{ normality of HCL} \times 14 \times 100 \times 100}{\text{Weight of the sample} \times 10 \times 1000} \dots\dots\dots 1\% \text{ protein} = \% \text{ N2} \times 6.25$$

Fat content: The fat content was determined with the help of Soxhlet apparatus as per the standard method (AOAC, 1995 [20]) and results were expressed in terms of percentage. Fat contribute to the flavor of food as well as it gives texture and also mouth feel to the food. It give us maximum energy 9Kcal energy per gram. To measure the amount of fat present in the food because extra intake of fat mostly leads to obesity and below the level lead to malnutrition. Soxtronis a solvent extraction method is more pronouncedly known as Soxhlet method.

Calculation: Fat (%) = (W2-W1) / W ×100

Where,
 W2 = Weight of flask with oil.
 W1 = Weight of empty flask.
 W= Weight of initial sample.

Fiber content: About 5 g of moisture and fat free sample was boiled in 500 ml beaker containing 200 ml boiling 0.255 N (1.25 w/v) H2SO4. The mixture was boiled for 30 min keeping the volume constant by addition of water at frequent intervals. At the end of this period, the mixture was filtered through a filter paper and the residue was washed with hot water till free from acid. The material then transferred to the same beaker and 200 ml of boiling 0.313 N NaOH solution added. After boiling for 30 min, the mixture was filtered through filter paper. The residue was washed with hot water till free from alkali. It was then transferred to crucible and dried in oven for overnight at 80-100 °C and weighed. The crucibles were heated in a muffle furnace at 550-600 °C for 4-5 hrs then cooled and weighed again. The difference in the weights represented the weight of the crude fiber (AOAC).

Results

The results obtained in the present investigation entitled “Development of cookies enriched with protein and fiber” summarized here with and discussed for their significance. The whole data obtained on various aspects of study is categorized under the following headings and subheadings. Effect of level of addition of Soya Flour and Banana Peel Powder on physicochemical properties of cookies

Table 1: Shows Parameters and Level of soya flour and BPP

Parameters	Level of soya flour and BPP				
	Control	T1	T2	T3	T4
Moisture (%)	3.6	3.8	4.0	4.1	4.3
Carbohydrate (%)	68.9	68.1	68.0	67.8	67.6
Protein (%)	7.26	8.06	9.01	10.35	10.59
Fat (%)	21.61	21.59	21.64	21.42	21.40
Fiber (%)	0.12	0.78	1.52	2.49	3.21
Ash (%)	0.64	0.71	0.76	0.98	1.02

Proximate composition of cookies

- **Moisture Content:** Moisture content is the amount of moisture present in given food sample. When the food is baked the moisture present in food is evaporated increasing keeping quality and reducing moisture levels. Cookies should have less than 5% moisture so that it appears crispy. In baking there is loss of moisture. From the above table it represents the moisture content at different levels of Soya flour and BPP. Control sample has 3.6% moisture content. Moisture content increase with increase in replacement of flour. Moisture content is lowest in the control sample as compared to other samples.
- **Estimation of carbohydrate:** As shown in the fig and table below the carbohydrate of prepared sample is found to be higher in control (68.9%) followed by other, in sample T1 (68.1%), sample T2 (68.0%), sample T3 (67.8%) and sample T4 (67.6%). Carbohydrate is higher in control as compared to other samples.
- **Protein content:** Protein content is nothing but amino acids present in sample. Control sample has 7.26% protein content. From table 1 it is shown that protein content increases with the addition of soya flour. Protein content is highest in biscuits prepared by adding soya flour at levels of 20% followed by 15%, 10%, 5%. Protein content is found out by finding out nitrogen content and then multiplying by its factor 6.25. Protein content of T3 was 10.35%.
- **Fat content:** Cookies are made with “shortening”. The fat was rubbed into the flour. The molecules of fat surround the flour particles and exclude water. This prevents the development of gluten in the dough. The fat is said to shorten the dough. Any increase in water in the mixture will tend to encourage development of gluten, which will make biscuits hard. Fat content was calculated by Soxhlet extraction method. 5 g samples were mixed with about 90 ml of n-hexane. The mixture was vigorously shaken with the separation flask knob opened at intervals to release the accumulated air pressure, which may burst the flask if left there. The fat in spirit was evaporated to dryness over a Soxhlet extraction, which extracts n-hexane from its solution of fat. Control sample has 21.61% fat. From the table 4.1 it is shown that fat content increases with increase in levels of addition of Soya flour. Fat content is highest in cookies with 20% replacement followed by 20, 15, 10, 5% fat content respectively.
- **Fiber Content:** Fiber content is highest in cookies prepared by adding 12% of BPP (T4) followed by 9%, 6% and 3%. Fiber content increased with addition of BPP.
- This study shows that BPP addition in cookies is accepted to certain levels beyond it damages the overall acceptability of product. T3 found to be 2.49% fiber.
- **Ash content:** Ash content refers to the amount of minerals present in the food. As the level of incorporation of BPP flour increases the ash content increases. The main reason of increment of ash content with increase of BPP is the fibers are rich in minerals. Ash content was determined by using muffle furnace. Sample in crucible were kept in furnace at 500-550 °C for 5-6 hrs and cooled in desiccators and calculated. It

found that Ash content was maximum in T4 sample followed by T3, T2, T1 and control sample.

- **Microbial Analysis:** Cookies are classified under low moisture food categories which has shelf of more than 3 months and could be extended to year. Due to low Availability of moisture cookies are not prone to bacterial growth easily but sometimes fungal infection can occur. Fungal colonies (CFU/gm) was calculated for control and selected sample T3.

Table 2: Microbial count in the samples

Group	Control	Sample T3	Permissible
Fungal count (CFU/gm)	11	22	50

Dilution factor = 1×10^3

Results

Summary: The experiment was carried out with the different level of Soya Flour and BPP to check the increase in nutritional level of cookies. The cookies were prepared by adding Soya flour at the levels of 5%, 10%, 15%, 20% and Banana Peel Powder at 3%, 6%, 9% and 12% respectively. Cookies were evaluated for its various physicochemical, nutritional and organoleptic quality parameters, and then study revealed that biscuits prepared by adding Soya flour at level 15% and 9% BPP gave a better acceptable taste, texture and flavor.

There was gradual increase in medicinal properties and organoleptic tests gave best results for level of T3 addition of samples. Cookies enriched with soya flour and BPP are rich in protein and fiber, antioxidants along with certain medicinal properties.

References

1. AACC. Approved methods of the American Association of Cereal chemists Methods. 10th Ed. St Paul, MN: AACC; c2000.
2. AOAC. Official methods of analysis of the association of official analytical chemist (18th ed.). Washington DC USA; c2006.
3. Akubor PI. Functional properties and performance of cowpea/plantain/wheat flour blends in biscuits PI ant Foods for Human Nutrition. 2003;58:1-8.
4. Norhidayah *et al.* Textural and sensorial properties of cookies prepared by partial substitution of wheat flour with unripe banana. IFRJ. 2014;21(6):2133-2139
5. Swati ShuklaSoybean (*Glycine max* L.) & its health benefits International Journal of Food Science and Nutrition ISSN: 2455-4898; c2014.
6. Ligia Portugal Gomes RebelloFlour of banana (Musa AAA) peel as a source of antioxidant phenolic compounds Food Research International. 2014;55:397-403.
7. Kulthe. Development of high protein and low calorie cookies J Food Sci Technol; c2011. DOI: 10.1007/s13197-011-0465-2
8. Anant S Kulkarni. Studies on development of high protein cookiesP-ISSN: 2349-8528 E-ISSN: 2321-4902
9. Anhwange B. A Chemical Composition of Musa sepientum (Banana) Peels JEAFC ISSN: 15794377
10. Sindhu Hruyia L. IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT) e-ISSN: 2319-2402, p-ISSN: 2319-2399. 2016;10(3):27-40.

11. Mohammad A. Phytochemicals and nutritional health benefits of soyplant International Journal of Nutrition, Pharmacology, Neurological Diseases; c2013, 3(1).
12. Hernawati *et al.* J Phys.: Conf. Ser. 812 012118; c2017.
13. Orhevba BA, Salaudeen HO. Optimisation of the production process of bread fortified with soya protein isolate and fonio flour blend. Int. J. Agric. Food Sci. 2022;4(2):26-36. DOI: 10.33545/2664844X.2022.v4.i2a.95
14. Sharma P. ICAR-IISR Soya bean production of kharif.
15. Sylvia Bursens Soya, Human Nutrition and Health; c2017. DOI: 10.5772/21951.
16. NeelaBolla K. Soybean Consumption and Health Benefits International; c2017.
17. Maria A. Formulation and Physicochemical and Sensorial Evaluation of Biscuit-Type Cookies Supplemented with Fruit Powders Plant Foods Hum Nutr. 2009;64:153-159. DOI: 10.1007/s11130-009-0118-z