



E-ISSN: 2706-8927
P-ISSN: 2706-8919
IJAAS 2020; 2(3): 98-103
Received: 11-05-2020
Accepted: 13-06-2020

Shail Bajpai
Ph.D. Thesis & Department of
Zoology, S.N. Sen P.G.
College, Kanpur, Uttar
Pradesh, India

BS Chandel
Ph.D., D.Sc., Biopesticides and
Toxicological Lab., P.G.
Department of Zoology,
D.B.S. College, Kanpur, Uttar
Pradesh, India

Corresponding Author:
Shail Bajpai
Ph.D. Thesis & Department of
Zoology, S.N. Sen P.G.
College, Kanpur, Uttar
Pradesh, India

An assessment of ecofriendly insecticidal toxicity of *Ocimum sanctum*, *Croton tiglium* and *Zingiber officinale* extractives against Bihar hairy caterpillar, *Spilarctia obliqua* Walker (Lepidoptera: Arctiidae)

Shail Bajpai and BS Chandel

Abstract

The Bihar hairy caterpillar, *Diacrisia (Spilarctia) obliqua* Walker are sporadic in nature has been in regular occurrence in northern Indo-Gangatic region, causing considerable damage to cruciferous crops and vegetables in our country. An experiment was conducted to test the insecticidal efficacy of six naturally occurring plant extracts against third instar larvae of *Diacrisia (Spilarctia) obliqua* Walker under the laboratory conditions. The bio-efficacy of six plant materials viz., *Croton tiglium* Linn., *Datura alba* Linn, *Ocimum santum* Linn, *Spilanthes acmella* Murr, *Thavetia nerfolia* Linn. and *Zingiber officinale* Linn. were prepared with the help of the Soxhlet apparatus under the laboratory conditions and one untreated control used for detail study. The laboratory trial was done in three glass petridishes (10cm diameter and replicated thrice by using each selected extracts per petridish. To record the mortality of ten larvae of *Spilarctia obliqua* were released inside each pair of petridishes and allow them to remain for 24 hours. These petridishes were kept as such under control conditions and mortality count was taken after 6, 12, 24 hours of exposure. The result observed that *O. santum* registered highest mortality (65.61%) to the 3rd instars larvae of *Spilarctia obliqua* Walk. when compared to other plant extracts as; *C. tiglium* (57.39) > *Z. officinale* (55.84) > *S. Acmella* (51.70) > *T. Nerfolia* (51.56) > *D. Alba* (37.93), respectively. The aforesaid results have focused on natural plant products as alternatives for insect-pest control.

Keywords: *Spilarctia obliqua*, *Ocimum santum*, *Croton tiglium* and caterpillars

Introduction

Provision of food has always been a challenging when hungeriness facing mankind. A major cornerstone in this challenge is the competition from insects which claim to be responsible for destroying one-fifth of the world's total crop production annually (Wyckhuys *et al.* 2013) [1]. India is basically an agro-based country and more than 80% of Indian population still depends by agricultural production. It is unanimously agreed that insect-pests are the main factors causing damage to crops adversely affects agricultural production. The monetary loss due to feeding by larvae and adult insects alone contributes to billion dollars per annum (Nelson *et al.* 2016) [2].

The Bihar hairy caterpillars, *Diacrisia obliqua* (Walk.) are sporadic in nature has been in regular occurrence in northern India, causing considerable damage to cruciferous crops and vegetables in our country (Chandel *et al.* 2004 and Tandon *et al.* 2004) [3, 4]. The pest is distractive in its larval stages. The damaged plant shows stunted growth and results in deterioration of yield. Indiscriminate use of synthetic insecticides has led to problems has been raised about the ill effects to human health and affecting environment and increase the costs of insect control (Chandel *et al.* 2018) [5].

In the quest of recent approaches to insect-pests control viz., discovery of cheap, biodegradable, least adverse effective and ecofriendly effective pesticides, the insecticides suit well as far as Indian conditions are concerned, where the mass of farmers lack resources and technical knowledge. It is required that production of insecticides of plants origin should be encouraged in our country (Chaudhay and Dadheech, 1989, Konar and Rai, 1990, Schmidt and Streloke, 1994, Tripathi *et al.* 2000, Dubey *et al.* 2004, Isman, 2006, Bajpai and Chandel, 2009, Dang *et al.* 2010; Kim *et al.*, 2011 and Valsalan and Gokuldas, 2015 [6, 7, 8, 9, 10, 11, 12, 13, 14, 15]. Majority of Indians are vegetarian and vegetables play an important role in

human diet, supplying some of nutrients, which are important source of mineral elements and vitamins. Family cruciferae provides some valuable vegetable and oil yielding crop like mustard, *Brassica campestris*, *Brassica napus*, *Brassica juncea*, radish, *Ravenous sativus*, cabbage, *Brassica oleracea* var. *capitata* etc. particularly in all places. A large number of insect-pests as, cabbage butterfly, *Pieris brassicae* Linn., tobacco caterpillar, *Spodoptera litura* Fabr., Bihar hairy caterpillar, *Spilarctia obliqua* Walk., cabbage borer, *Hellula undalis* Fabr., mustard sawfly, *Athalia proxima* Klug., cabbage semilooper, *Trichoplusia ni* Hub. and cabbage leaf webber *Crociodolomia binotalis* Zell. Are limiting factors of crops (Bharti and Chandel, 1017)^[16].

Therefore, the present investigation was aimed to understand the differential bioefficacy of the certain indigenous plant extract against polyphagous insect, Bihar hairy caterpillar, *Spilarctia obliqua* as this information will help to develop a sowed insect pest management programmed.

2. Materials and Method

2.1 Plant collection and Preparation of Powder: The plant materials of selected plants were collected in November 2008 from the cultivated field of farmers and wild areas in the vicinity of Kanpur region of Uttar Pradesh, India. The plant species was authenticated, and plant parts were dried under shade and then powdered using a mechanical grinder into a coarse powder. The selected plant materials were collected from the vicinity of Kanpur region during spring season. The identification of these collected leaves was confirmed by the plant taxonomist of Department of Botany D.B.S. College, affiliated to C.S.J.M. University, Kanpur.

2.2 Preparation of Extract and their Formulations: The powdered plant material was extracted with 80% methanol

using Soxhlet extraction apparatus. The solvent was removed completely under reduced pressure and a semisolid mass was obtained. The extracts were stored in a vacuum desiccator for further use (Schmutterer, 1988)^[17].

2.3 Rearing of Insect: Adult male and female of the Bihar hairy caterpillar *Diacrisia obliqua* (Walk.) were used for the present investigation. They were housed in a clean polypropylene cage and maintained under standard laboratory conditions (temperature 25±2 °C). They were fed natural diet. All experimental procedures described by Nagarkatui and Prakash, 1974 and Chandel and Sengar, 2017^[18, 19]

3. Experimental Protocol

Insecticidal test was carried out under laboratory condition against 3rd instars larvae of *Diacrisia obliqua*. The mustard leaves were used as food for the larvae of *Diacrisia obliqua*. Mustard leaves were treated with different concentrations (0.25, 0.5, 1.0, 1.5 and 2.0 percent) for two minutes. The treated leaves were left under electric fan for about half an hour, to make a dry film of the extracts on the leaves for each set of extract and one control. The treated foods were kept in jar (23cm x 10cm) on moist filter paper. The untreated leaves were dipped in Benzene +emulsified water only. Ten starved larvae of *Diacrisia obliqua* were released in each jar along with control. Three replicates per treatments were maintained. Number of larvae of *Diacrisia obliqua* died or moribund in all treatments and replications were recorded.

4. Statistical Analysis

The experimental results were expressed as mean ± standard error of mean (SEM). Statistical significance was analyzed by analysis of variance and P values of <0.001 were considered to be statistically significant (Abbott, 1926)^[20].

Table 1: *In vitro* effect of the different extractives for their mean mortality against *D. obliqua* irrespective of concentration and periods

Treatment (Plant extracts)	Con. (%)	Mean Mortality Percent After					
		6hrs.		12hrs.		24hrs.	
		T ₁	T.B.V. ₁	T ₂	T.B.V. ₂	T ₃	T.B.V. ₃
<i>Croton tiglium</i> Linn.	0.5	0.5	37.14	36.5	53.07	63.9	67.84
<i>Croton tiglium</i> Linn.	1.0	43.07	46.6	59.00	73.5	72.78	91.2
<i>Croton tiglium</i> Linn.	2.0	48.93	56.8	61.92	77.8	72.78	91.2
<i>Datura alba</i> Linn	0.5	26.64	21.4	32.08	28.2	38.59	38.9
<i>Datura alba</i> Linn	0.1	32.20	28.4	37.35	36.8	37.75	54.1
<i>Datura alba</i> Linn	2.0	36.87	36.0	39.35	40.2	50.07	50.8
<i>Ocimum santum</i> Linn.	0.5	48.93	56.8	57.00	70.3	63.93	80.7
<i>Ocimum santum</i> Linn.	1.0	54.78	66.7	61.92	77.8	75.00	93.3
<i>Ocimum santum</i> Linn.	2.0	61.92	77.8	81.14	97.6	83.84	98.9
<i>Spilanthes acmella</i> Murr.	0.5	35.00	32.9	46.92	53.4	50.85	60.1
<i>Spilanthes acmella</i> Murr.	1.0	46.92	53.4	52.77	63.4	59.21	73.8
<i>Spilanthes acmella</i> Murr.	2.0	39.14	39.94	45.00	50.0	50.85	60.1
<i>Thavetia nerfolia</i> Linn.	0.5	0.5	39.14	39.9	45.00	50.0	50.85
<i>Thavetia nerfolia</i> Linn.	1.0	45.00	50.0	48.93	56.8	59.00	70.5
<i>Thavetia nerfolia</i> Linn.	2.0	48.93	56.8	44.78	66.7	72.29	90.8
<i>Zingiber officinale</i> Linn.	0.5	45.00	50.0	52.77	63.4	54.78	66.7
<i>Zingiber officinale</i> Linn.	1.0	52.77	63.4	64.07	64.9	61.92	77.8
<i>Zingiber officinale</i> Linn.	2.0	54.78	66.7	59.00	73.5	70.07	88.4
Control	0.00	0.00	00.0	0.00	00.0	21.14	7.04

(T₁, T₂, T₃ = Treatments and TBV.₁, TBV.₂, TBV.₃ = Transformed Back Values)

C.D. for the treatment combination means = 0.147

i) C.D. for control vs. treated = 5.0499

ii) C.D. for insecticide means = 4.3879

- iii) C.D. for concentration means = 2.0973
- iv) C.D. for period means = 2.0903
- v) C.D. for insecticide and concentration means at the same period = 7.1368

The analysis of variance in table 1 shows that the main effect of insecticides, concentration, and periods as well as ‘control versus treated’ in first order and periods, concentrations in second order interactions are more highly

significant except the first order interaction “Insecticides X concentrations” and the second order which is non-significant. The effect of “Control versus treatment” is also significant at 0.5% level of significance.

Table 2a: Mean mortality percentage of *Diacrisia obliqua* in case of different combination under laboratory conditions

Treatment (Plant extracts)	Con. (%)	Mean Mortality percent After					
		6 hrs.		12 hrs.		24 hrs.	
		T ₁	T.B.V. ₁	T ₂	T.B.V. ₂	T ₃	T.B.V. ₃
<i>Croton tiglium</i> Linn.	0.5	37.14	36.5	53.07	63.9	67.84	85.8
<i>Datura alba</i> Linn	0.5	26.64	21.4	32.08	28.2	38.59	38.9
<i>Ocimum santum</i> Linn.	0.5	48.93	56.8	57.00	70.3	63.93	80.7
<i>Spilanthes acmella</i> Murr.	0.5	35.00	32.9	46.92	53.4	50.85	60.1
<i>Thavetia nerfolia</i> Linn..	0.5	39.14	39.9	45.00	50.0	50.85	60.1
<i>Zingiber officinale</i> Linn.	0.5	45.00	50.0	52.77	63.4	54.78	66.7
Control	0.00	0.00	00.0	0.00	00.0	21.14	7.04

(T₁, T₂, T₃ = Treatments and TBV.₁, TBV.₂, TBV.₃= Transformed Back Values)

Table 2b: Mean mortality percentage of *Diacrisia obliqua* in case of different combination under laboratory conditions

Treatment (Plant extracts)	Con. (%)	Mean Mortality percent After					
		6hrs.		12hrs.		24hrs.	
		T ₁	T.B.V. ₁	T ₂	T.B.V. ₂	T ₃	T.B.V. ₃
<i>Croton tiglium</i> Linn.	1.0	43.07	46.6	59.00	73.5	72.78	91.2
<i>Datura alba</i> Linn	1.0	32.20	28.4	37.35	36.8	37.75	54.1
<i>Ocimum santum</i> Linn.	1.0	54.78	66.7	61.92	77.8	75.00	93.3
<i>Spilanthes acmella</i> Murr.	1.0	46.92	53.4	52.77	63.4	59.21	73.8
<i>Thavetia nerfolia</i> Linn..	1.0	45.00	50.0	48.93	56.8	59.00	70.5
<i>Zingiber officinale</i> Linn.	1.0	52.77	63.4	64.07	64.9	61.92	77.8
Control	0.00	0.00	00.0	0.00	00.0	21.14	7.04

(T₁, T₂, T₃ = Treatments and TBV.₁, TBV.₂, TBV.₃= Transformed Back Values)

Table 2c: Mean mortality percentage of *Diacrisia obliqua* in case of different combination under laboratory conditions

Treatment (Plant extracts)	Con. (%)	Mean Mortality percent After					
		6 hrs.		12 hrs.		24 hrs.	
		T ₁	T.B.V. ₁	T ₂	T.B.V. ₂	T ₃	T.B.V. ₃
<i>Croton tiglium</i> Linn.	2.0	48.93	56.8	61.92	77.8	72.78	91.2
<i>Datura alba</i> Linn	2.0	36.87	36.0	39.35	40.2	50.07	50.8
<i>Ocimum santum</i> Linn.	2.0	61.92	77.8	81.14	97.6	83.84	98.9
<i>Spilanthes acmella</i> Murr.	2.0	39.14	39.94	45.00	50.0	50.85	60.1
<i>Thavetia nerfolia</i> Linn..	2.0	48.93	56.8	44.78	66.7	72.29	90.8
<i>Zingiber officinale</i> Linn.	2.0	54.78	66.7	59.00	73.5	70.07	88.4
Control	0.00	0.00	00.0	0.00	00.0	21.14	7.04

(T₁, T₂, T₃ = Treatments and TBV.₁, TBV.₂, TBV.₃= Transformed Back Values)

Table 3: Mean mortality % of *D. obliqua* larvae in different periods irrespective of treatment

Treatments	6hrs	TBV-1	12 hrs	TBV-2	72 hrs	TBV-3	GT	TBV-G
<i>C. tiglium</i>	43.04	46.6	57.99	71.9	71.14	89.6	57.39	71.0
<i>D. alba</i>	32.08	28.2	36.27	35.0	43.40	50.8	37.93	38.0
<i>O. santum</i>	55.21	67.5	66.68	84.5	74.00	92.5	65.31	82.5
<i>S. acmella</i>	44.89	49.8	52.23	62.5	57.99	71.9	51.70	61.6
<i>T. nerfolia</i>	44.89	48.9	49.97	58.6	60.38	75.6	51.56	61.4
<i>Z. officinale</i>	50.85	60.1	54.44	67.0	62.25	78.3	55.84	68.9
Control	0.00	0.00	00.0	0.00	21.14	7.04	7.04	1.5

(Figures in parenthesis represent mean percentage transformed back values.)

- C.D. for period means at the same insecticide = 5.0497
- C.D. for insecticide means at the same period = 5.1452

The table 3 and Figure 3, 3a and 3b reveals that the insecticide, *Ocimum santum* Linn leaves extract gave the best results when compared to other plant extracted insecticides. It killed 63.31 per cent of larvae followed by

Croton tiglium Linn. (57.39 per cent), *Zingiber officinale* Linn. (55.84 per cent), *Spilanthus acmella* Murr. (51.70 percent), *Thavetia nerfolia* Linn. (51.56 per cent) and *Datura Alba Linn.* (37.93 per cent), respectively. The

insecticide *Ocimum sanctum* differs significantly from the remaining ones except *Croton tiglium* from which it does not differ significantly to one another. *Datura Alba* proved

least toxic giving only 37.73 percent mortality of *Dicricia obliqua* Walker larvae.

Table 4: Mean mortality % of *D. obliqua* in concentrations irrespective of periods under the laboratory condition

Treatments extractivess	Mean mortality percent after						Mean % Mortality	
	6 hrs.		12 hrs.		24 hrs.		G.T.	TBV
	T ₁	TBV ₁	T ₂	TBV ₂	T ₃	TBV ₃		
0.5	41.08	43.2	46.38	52.4	57.63	71.3	48.36	55.9
1.0	47.65	54.6	55.67	68.2	66.49	84.1	56.60	69.7
2.0	57.63	47.3	62.52	78.7	76.45	94.5	66.53	82.8
Control	0.00	0.00	00.0	0.00	21.14	7.04	7.04	1.5

(T₁, T₂, T₃ = Treatments and T.B.V.₁, T.B.V.₂, T.B.V.₃ = Transformed Back Values)

Table 4 and figure indicate that all the three concentration differed significantly to one another. The concentration 2.0 per cent is superior to concentration 1.0 and 0.5 per cent. It is observe that the difference in the percentage larvae of *Dicricia obliqua* kill in concentration 2.0 percent and 1.0 per

cent is greater than the difference in concentration to kill the grub and adults in 1.0 per cent and 0.5 per cent in all the three periods. Similarly, the difference in percentage mortality of larvae of *Dicricia obliqua* Walker.

Table 5: Mean mortality % of *D. obliqua* in control irrespective of treatments.

Treatments extractivess	Mean mortality percent after						Mean % Mortality	
	6 hrs.		12 hrs.		24 hrs.		G.T.	TBV
	T ₁	TBV ₁	T ₂	TBV ₂	T ₃	TBV ₃		
Control	0.00	0.00	00.0	0.00	21.14	7.04	7.04	1.5
Extractives	16.18	7.8	18.83	10.4	21.82	13.8	18.95	10.5
Mean %	0.09	0.00	9.41	2.7	21.48	13.4	12.39	4.6

5. Result and Discussion

In the present study, administration of *Dicricia obliqua* larvae infested on mustard and vegetables can be controlled by selected biorational products. The leaves extract of *Ocimum santum* (Linn.) gave the best results when compared to other plant extracted insecticides. It killed 63.31 per cent of larvae followed by *Croton tiglium* Linn. (57.39 per cent), and *Zingiber officinale* Linn. (55.84 per cent), respectively. The insecticide *Ocimum santum* Linn. differs significantly from the remaining ones except *Croton tiglium* from which it does not differ significantly to one another. It is evident from the table A that the highest mortality of *D. obliqua* was observed 65.61 per cent treated with *Ocimum sanctum* concentration followed by *Croton tiglium* and *Zingiber officinale* giving 57.39 per cent and 55.84 percent mortality, within 24hrs. of treatment. Mortality percentage in all plant extracts treatment data were found highly significant as compared to control. The significance difference was observed in *Alpinia galanga* and *Zingiber officinale* treatment as compared to remaining plant extracts but no difference was observed. The higher mortality 65.61 per cent was recorded against third instars larvae of *D. obliqua* in case of *Ocimum sanctum* followed by *Croton tiglium* (57.39 per cent) and *Zingiber officinalis* (55, 84 per cent) as compared to control.

In the support of our findings many workers reported time to time that some plants protected themselves against injurious insects by producing their own biochemical defenses that are insecticidal bivalents (Bharti and Chandel, 2017, Chandel and Singh, 2017) [21, 22]. The consideration for the use of extracts of plants origin is that they are easily biodegradable, effective on some pests and considered safe in pest control operations as they minimize pesticide residues, ensure safety of the consumers of the treated grains and the environment (Thara and Kingsly, 2001, Gowri et al. 2002, Tomova et al. 2005) [23, 24, 25].

Further, the production of organic extracts of plant origin for pest control may be easier and less expensive than the synthesis of some complex chemical (Zygodlo et al. 1990, Olfa et al. 2009) [26, 27]. They possess many of the attributes of an ideal biological control agent, including broad host range, high virulence, host seeking capability, ease of mass production, recycling ability, non-hazardous to environment, etc. (Mahla et al. 2002, Sumathi et al. 2002, Panickar et al. 2003 and Ebrahimi et al. 2005) [28, 29, 30, 31]. Mansour et al. (2014) screened the toxicity of chicory, *Cichorium intybus* L. (Asteraceae), against larvae and adults of the mosquito (*Anopheles pharoensis*) and the housefly management. [32].

It can be concluded from above results that rhizome extracts of *Ocimum sanctum* and *Zingiber officinale* were most effective and gave highest mortality of third instars larvae of *Ocimum sanctum*, among all the selected plant extracts tested in the laboratory. Though, all the plant products were significantly better than control as regards the mortality of *S. obliqua* on mustard leaves are concerned after 6hrs, 12hrs. And 24hrs. Of treatments under laboratory conditions.

6. Conclusion

Conclusively, our findings reveal that the plant extract of *Ocimum sanctum* gave the maximum mortality. It killed 65.61 per cent larvae of *Spilarctia obliqua* followed by *Croton tiglium* (57.39 per cent) and *Z. officinale* (55.84 per cent) and control, respectively. The plant extract of *Ocimum sanctum* differed significantly from remaining once except *Croton tiglium* and *Z. officinale* from which it does not differs significantly to one another. Finally, it can be concluded on the basis of mean mortality percentage of six selected plant extracts against *S. obliqua* Linn. *D. Alba* is proved the least effective compare than all other plant extract. No physical injury was noticed on any part of the

plant after spraying. During the spraying of the extract it was noticed that extracts gave a strong irritating and unpleasant odor. Use of these selected plant extractives can be used for insect pest management.

7. Acknowledgement

The authors thank the Principal D.B.S. College, Kanpur and Head, Department of Zoology, D.B.S. College, Kanpur, India, for providing the necessary facilities to perform the present study.. Prof. R.A. Tripathi, farmer, Head, Division of Entomology, C S Azad University of Agriculture and Technology, Kanpur for rendering their support and help for the completion of this work and manuscript.

References

- Wyckhuys, Kris AG, Yanhui Lu, Helda Morales, Luis L. Vazquez, Jesusa C. Legaspi, Panagiotis A. Eliopoulos, Luis M. Hernandez. Current status and potential of the conservation biological control for agriculture in the developing world. *Biological control*. 2013; 65(1):152-167.
- Nelson Mpumi, Kelvin Mtei, Revocatus Machunda, Patrick A. Ndakidemi The Toxicity, Persistence and Mode of Actions of Selected Botanical Pesticides in Africa against Insect Pests in Common Beans, *P. vulgaris*: A Review, *American Journal of Plant Sciences*. 2016; 7(1):12-19.
- Chandel BS, Srivastava SC, Dwivedi ND, Shukla S, Dubey A. Bioefficacy of certain asteraceous plants on growth, development, nutrition and reproduction of *Spilarctia obliqua* Walker, *Nat J Life Science*. 2004; 1(2):4577-460.
- Tandon S, Mittal AK, Kasana, Pant AK. "Effect of essential oil of *Elsholtzia densa*. On growth and reproduction of *Spilosoma obliqua* (Walker) *Indian Journal of Entomology*. 2004; 66(3):206-208.
- Chandel BS, Sudhir, Tripathi, Arti Singh. Laboratory bioassay of N-(P-methoxy Phenyl-2 hydroxy salicylimide and N-(p-chlorophenyl)-2 hydroxysalicylimide as repellent against pulse beetle, *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). *International Journal of Entomology Research*. 2018; 3(1):265-270.
- Chaudhary HR, Dadheech LN. Incidence of insects attacking okra and the avoidable losses caused by them. *Ann. Arid Zone*. 1989; 28(3-4):305-307.
- Konar A, Rai L. Efficacy of some insecticides against shoot and fruit borer *Earias vittella* Fab. and *Earias insulana* (Boisd.) of okra (*Abelmoschus esculentus* L. Moench). *Environ. Ecol*. 1990; 8(1):410-413.
- Schmidt GH, Streloke M. Effect of *Acorus calamus* Linn. (Araceae) oil and its main compound B-Asarone on *Prostephanus truncatus* Horn (Coleoptera: Bostrichidae) *J Stored Product Research*. 1994; 30(3):227-235.
- Tripathi AK, Prajapati V, Agarwal KK, Khanuja SPS, Kumar S. Repellency and toxicity of oil from *Artemisia annua* to certain stored product beetles. *Jour. Econ. Ent*, 2000; 93(1):43-47.
- Dubey A, Gupta R, Chandel BS. Efficacy of *Acorus calamus*, *Vitex negundo* and *Ageratum conyzoides* against tobacco caterpillar *Spilarctia obliqua* walker". *Indian. J Entomology*. 2004; 66(3):238-240.
- Isman MB. Botanical Insecticides, Deterrents, and Repellents in Modern Agriculture and an Increasingly Regulated World. *Annual Review of Entomology*. 2006; 51:45-66.
- Bajpai R, Chandel BS. Evaluation of *Adhatoda vasica*, *Acorus calamus* and *Vitex nungundo* as a source of phago-antifeedant against cabbage butterfly, *Pieris brassicae* Linn. (Lepidoptera: Pieridae) on mustard, *Brassica campestris* (Linn.). *International Journal of Biotechnology and Biochemistry*. 2009; 5(\$):451-459.
- Dang QL, Lee GY, Choi YH, Choi GJ, Jang KS, Park MS *et al*. Insecticidal activities of crude extracts and phospholipids from *Chenopodium ficifolium* against melon and cotton aphid, *Aphis gossypii*. *Crop Prot*. 2010; 29:1124-1129.
- Kim SK, Kim YC, Lee S, Kim JC, Yun MY, Kim IS. Insecticidal activity of rhamnolipid isolated from pseudomonas sp. EP-3 against green peach aphid. *J Agric Food Chem*. 2011; 59(3):934-938.
- Valsalan KK, Gokuldas M. Repellent and oviposition deterrent effects of *Clerodendrum infortunatum* on the pulse beetle *Callosobruchus chinensis* L. (Coleoptera: Bruchidae), *Journal of Entomology and Zoology Studies*. 2015; 3(4):250-255.
- Bharti, Anju, Chandel BS. Biorational and Ecofriendly Insecticidal Approach of Asteraceous Plant Extract against Spotted Bollworm, *Earias vittella* Fabricius (Lepidoptera: Noctuidae) on Okra, *Abelmoschus esculentus* Linn. (Moench) in Kanpur Region. *Journal of Entomology and Zoology Studies*. 2017; 5(4):1993-1999.
- Schmutterer H. Potential of azadirachtin-containing pesticides for integrated pest control in developing and industrial countries. *J Insect Physiol*. 1988; 34:713-719.
- Nagarkatti S, Prakash A. Rearing *Heliothis armigera* (Hubn) on artificial diet. *Commonwealth Institute of Biological Control, Bangalore, Technical Bulletin*. 1974; 17:169-173.
- Chandel BS, Sengar HS. Insectifuge effect of neem, *Azadirachta indica* A. Juss and their derivatives against sawfly, *Athalia lugens proxima* Klug. (Hymenoptera: Tenthredinidae). *International Journal of Zoology Studies*, 2018; 3(1):53-59.
- Abbott WS. A method of computing the effectiveness of insecticides. *Journal of Economic Entomology*. 1925; 18:265-267.
- Bharti, Anju, Chandel BS. Insectifuge biopotency of *Chrysanthemum cinerariifolium*, *Cichorium intybus*, *Inula racemosa* and *Tagetes minuta* extractives against okra aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae), *International Journal of Fauna and Biological Studies* 2017; 4(3):133-138.
- Chandel BS, Singh, Arti. Enterotoxicity of *Chromolaena odorata*, *Tagetes minuta* and *Reichardia tingitana* in suppressing oviposition and adult emergence of *Callosobruchus chinensis* (L) infesting stored chickpea seeds in U.P. *International Journal of Zoology Studies*. 2017; 2(6):38-44.
- Thara S, Kingsly S. Antifeedant effect of neem on bhindi pest *Earias vittella* (Lepidoptera: Noctuidae). *J Ecotoxicol. and Environ. Monito*. 2001; 11(2):131-134.
- Gowri S, Ramachandra Rao G, Nagalingam B. Evaluation of certain new neem formulations against

- okra fruit borer, *Earias vittella* (Fabricius) and their effect on yield. J Ento. Res. 2002; 26(3):245-247.
25. Tomova BS, Waterhouse JS, Doberski J. The effect of fractionated *Tagetes* oil volatiles on aphid reproduction. Entomologia Experimentalis et Applicata, 2005; 115:153-159.
26. Zygadlo JA, Grosso NR, Alburra RE, Guzman CA. Essential oil variation in *Tagetes minuta* populations. Biochem. Syst. Ecol. 1990; 18:405-407.
27. Olfa B, Monia BK, Samia A, Dalila H, Zine M, Ahmed NH. Insecticidal activity of some Asteraceae plant extracts against *Tribolium confusum*. Bulletin of Insectology. 2008; 61(2):283-289.
28. Mahla, Mukesh, Singh, Ram, Puja, Suhag *et al.* Biological efficacy of *Melia azedarach* roots against *Earias vittella* larvae. Journal of Medicinal and Aromatic Plant Sciences. 2002; 24(3):726-728.
29. Sumathi E, Balasubramanian G. Integrated management tactics for bhendi [*Abelmoschus esculentus* (L.) Moench] fruit borers, *Earias vittella* Fabricius and *Earias insulana* Boisduval. Res. On Crops. 2002; 3(1):171-176.
30. Panickar, Bindu, Bharpoda TM, Patel JR, Patel JJ. Evaluation of various schedules based on botanical and synthetic insecticides in okra ecology. Indian J of Ento. 2003; 65(3):344-346.
31. Ebrahimi M, Hasan Safaralizade M, Valizadegan O. Contact toxicity of *Azadirachta indica* (Adr. Juss.), *Eucalyptus camaldulensis* (Dehn.) and *Laurus nobilis* (L.) essential oils on mortality cotton aphids, *Aphis gossypii* Glover (Hemiptera: Aphididae). Archives of Phytopathology and Plant Protection, 2013; 46:2153-2162.
32. Masour SA, Ibrahim RM, Gengaihi SEEL. Insecticidal activity of chicory (*Cichorium intybus* L.) extracts against two dipterous insect-disease vectors: Mosquito and housefly, Industrial Crop Products. 2014; (254):192-202.